



Physico-Chemical and Sensory Evaluation of Chicken Nuggets with Cooked Unripe Plantain

K. Selvakumar, M. Anna Anandh*, G. Rajarajan, R. Ilavarasan and A. Sangeetha¹

Department of Livestock Products Technology, Tamil Nadu Veterinary and Animal Sciences University (TANUVAS), Veterinary College and Research Institute, Orathanadu- 614 625, Thanjavur District, Tamil Nadu, India.

¹Department of Veterinary Public Health and Epidemiology

ARTICLE INFO

* Corresponding author.

E-mail address: drannaanandh@gmail.com

Received 2023-12-07; Accepted 2024-03-28

Copyright © Indian Meat Science Association
(www.imsa.org.in)

DOI: 10.48165/jms.2024.19.01.07

ABSTRACT

Chicken nuggets were prepared with 5%, 10%, and 15% steam cooked unripe plantain without peel, and their qualities were evaluated. Linear and significantly ($P < 0.01$) increased values were observed for emulsion stability, cooking yield, moisture retention, fat retention, moisture, and total ash contents. Conversely significantly ($P < 0.01$) reverse trends were observed for emulsion pH, product pH, protein and fat contents. Fiber content values were significantly increased from 0.46 ± 0.03 to $5.71 \pm 0.08\%$. Sensory evaluation done using nine point hedonic scale indicated that 10% cooked unripe plantain without peel resulted in higher scores for appearance, colour, flavor, juiciness, tenderness, texture and overall acceptability among the three treatments. Thus, it can be concluded that, 10% cooked unripe plantain without peel can be successfully used for preparation of fiber enriched functional chicken nuggets of acceptable quality.

Key words: Acceptability, Banana, Chicken, Fiber, Functional, Meat, Nuggets, Quality, Plantain

INTRODUCTION

Meat is a nutritious food which can fulfill most of the physiological and functional requirements of human body mechanisms. Poultry meat is preferred by majority of the people in our country due to lower cost, higher availability and fact that chicken meat does not have any religious taboo (Talukder and Sharma, 2010). Chicken meat has high protein content, low fat content coupled with a balanced n – 6 to n – 3 polyunsaturated fatty acids ratio, low cholesterol and presence of some of the functional components (Petracci *et al.*, 2013). Despite these

nutritional advantages, chicken meat lacks dietary fiber, a crucial element for normal physiological and biochemical processes, emphasizing the significance of incorporating dietary fibers into meat products to support overall nutrition and digestive health. The inclusion of dietary fiber from diverse plant sources has the potential to mitigate the adverse effects associated with meat consumption while enhancing the nutritional composition and desirability of these products. Notably, dietary fiber's major functional properties, such as water holding capacity, gel-forming ability, and fat binding capacity, are of particular interest to meat processors (Kim *et al.*, 2012)

Fruits, recognized as an important source of dietary fiber, include bananas, which stand out as one of the most widely distributed and consumed fruits in tropical and subtropical regions. Bananas, easily accessible to the general population. It can be consumed as cooked as well as in uncooked forms (Kookal and Thimmaiah, 2018). Bananas are known for its health benefits for humans as it's packed with nutrients similar as potassium, calcium and iron. It's also good source of filaments and vitamins, bioactive composites such as phenolic composites, and resistant bounce potentially contributing to health benefits rendering it a fit seeker for classifying it as a functional food (Vu *et al.*, 2018). Recently, plantains have gained attention as a component for functional foods, particularly because of the low insipidity of plantain bounce. This property makes it an excellent component for food processing (Ming-Chang Li *et al.*, 2020). Unripe banana and banana by products are high in dietary fiber and phenolic compounds, and it has high antioxidant, antibacterial and antibiotic activities (Fidrianny and Insanu 2014; Pereira *et al.*, 2020). Thus, the incorporation of plantain components in various functional foods is of interest. However, use of unripe cooked banana and ripened banana products as a source of dietary fiber in meat products is yet to be fully explored. Hence, a scientific approach was made to develop functional chicken nuggets with unripe cooked plantain without peel.

MATERIALS AND METHODS

A. Raw Materials:

i. Chicken meat

Deboned chicken meat was purchased from the local meat stall. It was cut into small chunks and frozen for 1-2 hrs to ensure easy mincing. The chicken meat chunks were minced twice through the meat mincer and the minced meat was used in the preparation of chicken nuggets.

ii. Preparation of cooked unripe plantain

Monthan variety of plantain was used in this study. *Monthan* variety is exclusive culinary variety, widely cultivated for processing in Tamil Nadu. Fruits are large, peel is thick and pulp - firm, cream coloured, core conspicuous, medium taste. The mature unripe *Monthan* variety plantain of uniform size and shape were purchased from the local market. Before the plantain was cooked, the adhering extraneous materials on the surface were removed by using knife and washing in the running water. Then the plantain was steamed for 30 min to an internal temperature of 80°C. The cooked plantain was peeled. The cooked plantain without peel were then shredded using a vegetable shredder and then require quantity of shredded cooked plantain without peel was used in the chicken nuggets preparation based on the formula.

B. Formulation and Treatments

Table 1: Formulation of functional chicken nuggets

Ingredients	C - %	Level of cooked unripe plantain (%)		
		T1 - 5%	T2 - 10%	T3 - 15%
Chicken meat	73.0	68.0	63.0	58.0
Salt	1.5	1.5	1.5	1.5
Sodium tri poly phosphate	0.5	0.5	0.5	0.5
Dry spice mix	1.5	1.5	1.5	1.5
Condiments mix	3.0	3.0	3.0	3.0
Vegetable oil	7.5	7.5	7.5	7.5
Refined wheat flour	3.0	3.0	3.0	3.0
Ice flakes	10.0	10.0	10.0	10.0

The formula for chicken nuggets was developed after conducting a series of preliminary trails. Functional chicken nuggets were prepared with incorporation of 5, 10 and 15% cooked unripe plantain. The corresponding levels of chicken were 68%, 63% and 58% in the respective treatment. Other ingredients used in the formulation ie., common salt, sodium tri poly phosphate, dry spice mix (aniseed - 10, black pepper - 10, caraway seed - 10, capsicum - 8, cardamom - 5, cinnamon - 4, clove - 1, coriander - 20, cumin seed - 22, turmeric 10 in percentage/weight), condiments mix (onion and garlic in the ratio of 4:1), vegetable oil (Gold winner brand), refined wheat flour and ice flakes. Control chicken nuggets contained 73% chicken meat without cooked unripe plantain without peel (Table. 1).

C. Preparation of functional chicken nuggets

Chicken meat was partially thawed at refrigeration temperature (4±1°C) and the thawed meat was cut in to small cubes and minced in the meat mincer (Model: Primus MEW 713, Mado GmbH, Germany) using 8 mm plate. Pre weighed quantities of salt and sodium tripolyphosphate (STPP) were added to the minced chicken meat and chopped for 2 min with a food cutter (Model TC 11, Schaefer, Germany). After addition of ice flakes, it was chopped again for 1-2 min. Refined vegetable oil was incorporated slowly and chopping was continued till the oil and fat were completely dispersed in the batter. Condiments mix, spice mix and refined wheat flour as binder were added and chopping continued for 2 min to give a fine viscous emulsion. The temperature of the emulsion varied from 10 to 12°C. The emulsion was then filled uniformly in to aluminium moulds, packed compactly and covered. The moulds were then clipped and tied and the meat blocks were cooked in a steam oven without pressure for 40 min. The internal core temperature of the cooked blocks was 80 ± 2°C measured using a probe type thermometer (Acetec, India). The meat blocks were cooled to

room temperature, chilled overnight at 4 ± 1 °C and cut into slices of 15 mm thickness using a meat slicer (ALS, India). The slices were manually cut into nuggets (1 cm x 1 cm). The fried nuggets were served immediately to a panel for evaluation of various sensory attributes on a 9 – point hedonic scale.

D. Physico- Chemical Analysis

i. pH

The pH of the emulsion and product samples were determined by homogenizing 10 gram of sample with 50 ml distilled water with the help of tissue homogenizer for 1 minute. The pH of the suspension was recorded by immersing the combined glass electrode of digital pH meter (Model LMPH 9Labman, India). The pH meter was pre calibrated using standard solution with pH 7.0 as per the user manual instructions, prior to measurement.

ii. Emulsion stability (%)

The method was based on the emulsion stability test reported by Baliga and Madaiah (1971) with slight modifications. 20 gram of emulsion was rolled into balls and placed in polyethylene bags. The samples were cooked at 80°C in a thermostatically controlled water bath for 20 minutes. After draining out of the exudate, the cooked mass was cooled and weighed again. The emulsion stability was calculated as percentage by using the below mentioned formula.

$$\text{Emulsion stability (\%)} = \frac{\text{Weight of emulsion after cooking}}{\text{Weight of raw emulsion}} \times 100$$

iii. Product yield (%)

Product yield is used to predict the yield of cooked product. The product yield was determined by the method outlined by Anna Anandh *et al* (2008). The weight of the product was recorded before and after cooking from which the cooking yield was calculated by using the following formula.

$$\text{Product yield (\%)} = \frac{\text{Weight of product after cooking}}{\text{Weight of product before cooking}} \times 100$$

iv. Moisture retention

The moisture retention value represents the amount of moisture retained in the cooked product per 100 gm of sample and was determined according to El-Magoli *et al.* (1996) according to the equation below:

$$\text{Moisture retention (\%)} = \frac{\% \text{ yield} \times \% \text{ moisture in chicken nuggets}}{100} \times 100$$

v. Fat retention

The fat retention value represents the amount of fat retained in the product after cooking. Fat retention was calculated according to Murphy *et al.* (1975) by using the equation as follows:

$$\text{Fat retention (\%)} = \frac{(\text{Cooked weight}) \times (\text{Fat in cooked chicken nuggets})}{(\text{Uncooked weight}) \times (\text{Fat in uncooked chicken nuggets})} \times 100$$

vi. Proximate composition

The moisture, protein and fat contents of control and functional chicken nuggets were determined by standard methods using hot air oven, kjeldahl's assembly, soxhlet extraction apparatus and muffle furnace, respectively (AOAC, 1995). Estimation of the dietary fibre content of functional chicken meat nugget was done by using modified method prescribed by AOAC (2012).

E. Sensory Evaluation

Sensory evaluation was conducted with semi-trained panelists. Control and cooked unripe plantain incorporated chicken nuggets were served to the panelists. The sensory attributes like appearance and colour, flavour, juiciness, tenderness, texture and overall acceptability were evaluated on 9 - point descriptive scale (where in 1 is extremely undesirable and 9 is extremely desirable) as suggested by Keeton (1983).

F. Statistical Analysis

The experiment was repeated six times. The data generated from each trial were analyzed statistically by following standard procedures (Snedecor and Cochran, 1989) for comparing the means and to determine the effect of treatment.

RESULTS AND DISCUSSION

Physico- Chemical Characteristics

Results of physico-chemical parameters and proximate composition of chicken nuggets incorporated with different levels of cooked unripe plantain without peel (T1 - 5%, T2 - 10%, and T3 - 15%) and control (cooked unripe plantain without peel) are presented in Table are presented in Table 2.

The values for emulsion and product pH progressively and significantly ($P < 0.01$) increased from control to cooked unripe plantain without peel incorporated chicken nuggets. Significantly ($P < 0.01$) lower pH value was observed in T3 followed by T2, T1 and control.

This trend may be attributed to the lower pH of the added green banana pulp (Izidoro *et al.*, 2008), aligning with the results reported by Pereira *et al.* (2020) in fat-reduced frankfurters incorporated with unripe banana by-products. Carballo *et al.* (2021) also reported an increase in batter pH with an increase in banana pseudo-stem in pork patties.

The mean $\pm$ SE emulsion stability and product yield increased significantly from control to cooked unripe plantain without peel incorporated chicken nuggets. Among treatments, higher product yield observed in T3 followed by T2 and T1. This increase could be attributed to the fibers present in unripe banana by-products, allowing the retention of more water due to its absorption capacity (Pereira *et al.*, 2020). The emulsion stability is directly proportional to the cooking yield percentage, which is in agreement with Pereira *et al.* (2020), who observed increased cooking yield in fat-reduced frankfurters with the addition of banana by-products obtained from unripe fruit. Similar trends were observed in Bologna-type sausages formulated with green banana and pork skin gel (Alves *et al.*, 2016).

Table 2: Effect of incorporation of cooked shredded unripe plantain without peel on physico-chemical properties and proximate composition of chicken nuggets (Mean $\pm$ SE)

Parameters	Level of cooked unripe plantain without peel			
	C - 0%	T1 - 5%	T2 - 10%	T3- 15%
Physico-chemical parameters*				
Emulsion pH	6.22 $\pm$ 0.00 ^a	6.16 $\pm$ 0.00 ^b	6.12 $\pm$ 0.01 ^b	6.07 $\pm$ 0.01 ^c
Emulsion stability (%)	94.24 $\pm$ 0.53 ^a	95.50 $\pm$ 0.41 ^b	95.89 $\pm$ 0.36 ^b	96.37 $\pm$ 0.52 ^c
Cooking yield (%)	94.29 $\pm$ 0.78 ^a	95.49 $\pm$ 0.60 ^b	95.86 $\pm$ 0.36 ^b	96.32 $\pm$ 0.33 ^c
Product pH	6.31 $\pm$ 0.00 ^a	6.24 $\pm$ 0.00 ^b	6.19 $\pm$ 0.01 ^b	6.15 $\pm$ 0.00 ^c
Moisture retention (%)	57.99 $\pm$ 0.60 ^a	59.82 $\pm$ 0.88 ^b	61.20 $\pm$ 0.44 ^c	62.62 $\pm$ 0.51 ^d
Fat retention (%)	90.13 $\pm$ 1.48 ^a	91.04 $\pm$ 0.80 ^b	92.24 $\pm$ 0.51 ^b	92.99 $\pm$ 0.80 ^c
Proximate composition*				
Moisture(%)	61.51 $\pm$ 0.40 ^a	62.65 $\pm$ 0.70 ^b	63.85 $\pm$ 0.57 ^c	65.02 $\pm$ 0.48 ^d
Protein(%)	16.65 $\pm$ 0.50 ^a	16.46 $\pm$ 0.54 ^a	15.70 $\pm$ 0.16 ^b	14.68 $\pm$ 0.23 ^c
Fat(%)	20.57 $\pm$ 0.15 ^a	19.49 $\pm$ 0.13 ^b	18.55 $\pm$ 0.15 ^c	17.61 $\pm$ 0.13 ^d
Total ash(%)	2.52 $\pm$ 0.05 ^a	2.68 $\pm$ 0.05 ^b	2.89 $\pm$ 0.04 ^c	3.22 $\pm$ 0.21 ^d
Fiber content (%)	0.46 $\pm$ 0.03 ^a	2.15 $\pm$ 0.02 ^b	4.38 $\pm$ 0.02 ^c	5.71 $\pm$ 0.08 ^d

Number of observations: * = 6

Means $\pm$ SE bearing same superscripts row-wise do not differ significantly.

The mean $\pm$ SE moisture retention and fat retention values of control and cooked unripe plantain without peel incorporated chicken nuggets differed significantly

($P < 0.01$) between them and the value were significantly ($P < 0.01$) increasing with increasing level of cooked unripe plantain without peel in the formulation. Among cooked unripe plantain without peel incorporated chicken nuggets moisture retention and fat retention were higher in T3 followed by T2, T1 and control. Pereira *et al.* (2020) reported higher retention of moisture in unripe banana flour could be an efficient ingredient with the capacity to increase water retention. The increased water retention may be explained by the dietary fiber content of cooked unripe banana. Since dietary fiber is hydrated, the pore space in fiber particles is occupied by water molecules. Higher fat retention of cooked shredded unripe plantain without peel inclusion in chicken nuggets might be due to high total fiber content in the unripe green banana flour, which form gels at high temperatures, resulting in greater retention of fat in the products (Alves *et al.*, 2016).

Proximate composition

The mean $\pm$ SE moisture content and total ash content of T3 was significantly ($p < 0.01$) higher as compared to other treatments. Control had significantly ($p < 0.01$) lower moisture and total ash contents. This is in agreement with Pereira *et al.* (2020) who reported linear increase in moisture content in frankfurters formulated with increase in unripe banana by products. Higher values of ash content of cooked shredded unripe plantain without peel incorporated chicken nuggets might be due to higher concentration of total dietary fibers, resistant starch and minerals present in the added unripe plantain (Alves *et al.*, 2016).

The mean $\pm$ SE values of protein and fat content were significantly ($p < 0.01$) lower in T3 followed by T2, T1 and control. Control had significantly ($p < 0.01$) higher protein and fat contents. Similar results were reported by Troutt *et al.* (1992) in low fat beef patties incorporated with dietary fibers at different levels. The possible decrease in the crude protein content of cooked shredded unripe plantain without peel incorporated chicken nuggets might be due to replacement of lean meat with unripe plantain which contains low protein and high carbohydrate content. Choi *et al.* (2010) who successfully decreased the fat content by adding grape seed oil and rice bran fibers as a fat substitute. Santhi and Kalaikannan (2014) also observed significant decrease in protein and fat levels with increased levels of added vegetable fibers in chicken nuggets

The mean $\pm$ SE fiber content values were 0.46 $\pm$ 0.03, 2.15 $\pm$ 0.02, 4.38 $\pm$ 0.02 and 5.71 $\pm$ 0.08 for control, T1, T2 and 15%, respectively. The 15% cooked unripe plantain without peel incorporated chicken nuggets had significantly ($p < 0.01$) higher fiber content than the 10%, 5% cooked

unripe plantain without peel incorporated chicken nuggets and control had significantly ($p < 0.01$) lower among all formulation. Fiber content values between different treatments and the control differed significantly. This result aligns with Salazar *et al.* (2021), who observed higher fiber content in frankfurter sausages prepared with green banana products.

Sensory characteristics

Results of sensory evaluation of chicken nuggets incorporated with different levels of cooked unripe plantain without peel (T1 - 5%, T2 - 10%, and T3 - 15%) and control (cooked unripe plantain without peel) are presented in Table 2.

The mean $\pm$ SE appearance and colour scores significantly decreased from control to different level of cooked unripe plantain without peel inclusion in chicken nuggets. However, no significant difference observed between control, 5% and 10% level of inclusion and the values differed significantly from 15% level of inclusion. The mean scores $\pm$ SE for flavor, juiciness and texture were significantly ($P < 0.01$) higher for control as compared to cooked unripe plantain without peel incorporated chicken nuggets. Among treatments, the mean score for flavor, juiciness, tenderness and texture were higher for 10% cooked unripe plantain without peel incorporated chicken nuggets followed by 5% and 15% cooked unripe plantain without peel incorporated chicken nuggets. Mean score for overall acceptability was significantly higher for control followed by 10%, 5% and 15% cooked unripe plantain without peel incorporated chicken nuggets. No significant differences observed among the treatments but values was higher for T2, but the values were non significantly higher for 10% cooked unripe plantain without peel incorporated chicken nuggets than other two level of cooked unripe plantain without peel incorporated chicken nuggets. Akram *et al.* (2022) reported that colour scores of nuggets supplemented with banana peel powder were reduced because of the dark brown colour of plantain components. These results are in agreement with Pereira *et al.*, (2020), who reported incorporation of unripe banana by products in frankfurters formulation improved the sensory scores of the products. Garcia *et al.* (2002) reported that the incorporation of fiber-rich ingredients led to a decrease in the sensory scores of the sausage. A similar result of decreasing sensory scores with an increasing level of banana flour was also reported by Zaini *et al.* (2021) in chicken sausages. Zaini *et al.* (2020) reported that banana peel powder in concentrations of $>2\%$ decreased the organoleptic perception of sausage. The result of the

present study is in agreement with Akram *et al.*, (2022) who found that sensory evaluation of broiler nuggets made with banana peel powder decreased with an increasing level in the formulation.

Based on the results of sensory attributes, chicken nuggets prepared with 10% cooked unripe plantain without peel was rated better for all sensory attributes except appearance and colour. Tenderness score between treatments and control did not differed significantly. Among treatments, the best sensory response in terms of flavour, juiciness, tenderness, texture and overall acceptability was obtained for the chicken nuggets containing 10% cooked unripe plantain without peel. The result showed that cooked unripe plantain without peel could be successfully incorporated in the preparation of functional chicken nuggets even at 10% level without any adverse effects on the physico – chemical and proximate composition of chicken nuggets.

Table 3: Effect of incorporation of cooked shredded unripe plantain without peel on sensory attributes of chicken nuggets (Mean + SE)

Parameters	Level of cooked shredded unripe plantain without peel			
	C - 0%	T1 - 5%	T2 - 10%	T3- 15%
Sensory attributes**				
Appearance and colour	8.35 $\pm$ 0.28 ^a	8.27 $\pm$ 0.30 ^a	8.02 $\pm$ 0.30 ^a	7.52 $\pm$ 0.44 ^b
Flavor	8.02 $\pm$ 0.30 ^a	7.35 $\pm$ 0.48 ^b	7.72 $\pm$ 0.75 ^b	7.22 $\pm$ 0.63 ^b
Juiciness	8.12 $\pm$ 0.27 ^a	7.35 $\pm$ 0.23 ^b	7.50 $\pm$ 0.28 ^b	6.87 $\pm$ 0.22 ^c
Tenderness	7.55 $\pm$ 0.45	7.62 $\pm$ 0.53	7.80 $\pm$ 0.71	7.75 $\pm$ 0.59
Texture	8.20 $\pm$ 0.25 ^a	7.50 $\pm$ 0.51 ^b	7.45 $\pm$ 0.39 ^b	7.20 $\pm$ 0.52 ^b
Overall acceptability	8.27 $\pm$ 0.25 ^a	7.45 $\pm$ 0.42 ^b	7.65 $\pm$ 0.40 ^b	7.45 $\pm$ 0.42 ^b

Number of observations: ** = 20

Means $\pm$ SE bearing same superscripts row-wise do not differ significantly.

Sensory attributes of chicken nuggets were evaluated on a 9-point descriptive scale (wherein 1 = extremely undesirable; 9 = extremely desirable).

CONCLUSION.

Finding of the study suggests that incorporating cooked unripe plantain without peel in chicken nuggets has potential benefits in terms of nutritional enhancement, especially at the 10% level, without adversely affecting the overall sensory attributes. This research provides valuable insights into the development of functional meat products, considering both nutritional and sensory aspects.

REFERENCES

- Akram T, Mustafa S, Ilyas K, Tari MR, Al SW, Ali S, ShafiqM, Rao M, Safdar W, Iftikhar M, Hameed A, Manzoor M, Akhtar M, Umer Z and Basharat Z (2022) Supplementation of banana peel powder for the development of functional broiler nuggets. *Peer J.*, 10: e14364.
- Álvarez D and Barbut (2013) Effect of inulin, β -Glucan and their mixtures on emulsion stability, color and textural parameters of cooked meat batters, *Meat Sci.*, 94 (3):320-327.
- Alves, LAADS, Lorenzo JM, Gonçalves CA A, Alves dos Santos B, Heck RT, Cichoski AJ Campagnol PCB (2016) Production of healthier bologna type sausages using pork skin and green banana flour as a fat-replacers. *Meat Sci.*, 121: 73 -78.
- Anna Anandh M, Radha K, Lakshmanan V and Mendiratta SK (2008) Development and quality evaluation of cooked buffalo tripe rolls. *Meat Sci.*, 80: 1194–1199.
- AOAC (American Official Analytical Chemist) (1995) Official methods of analysis. 16th Edition, Virginia, USA.
- AOAC (American Official Analytical Chemist) (2012) Official methods of analysis. 19th Edition, Virginia, USA.
- Baliga BR and Madaiah N (1971) The quality of sausage emulsion prepared from mutton. *J. Food Sci.*, 35: 383.
- Carballo DE, Caro L, Gallego C, González AR, Giraldez FJ, Andres S and Mateo J (2021) Banana Pseudo-Stem Increases the Water-Holding Capacity of Minced Pork Batter and the Oxidative Stability of Pork Patties. *Foods*. 10: 2173.
- Choi YS, Choi JH, Han DJ, Kim HY, Lee MA, Kim HW, Lee JW, Chung HJ and Kim CJ (2010) Optimization of replacing pork back fat with grape seed oil and rice bran fiber for reduced-fat meat emulsion systems. *Meat Sci.*, 84(1): 212 – 218.
- El-Magoli SB, Laroia S and Hansen PMT (1996) Flavour and texture characteristics of low-fat ground beef patties formulated with whey protein concentrate. *Meat Sci.*, 42: 179 - 193.
- Fidrianny I. and InsanuM (2014) In vitro antioxidant activities from various extracts of banana peels using ABTS, DPPH assays and correlation with phenolic, flavonoid, carotenoid content. *Int. J Pharm and Pharmaceu. Sci.*, 6: 299 – 303.
- Garcia L, Dominguez R, Galvez D, Casas C and Selgas D (2002) Utilization of cereal and fruit fibres in low fat dry fermented sausages. *Meat Sci.*, 60: 227 – 232.
- Izidoro DR, Scheer AP, Sierakowski M and Haminiuk CWI (2008) Influence of green banana pulp on the rheological behaviour and chemical characteristics of emulsions (mayonnaises). *LWT Food Sci and Technol.*, 41: 1018 – 1028.
- Karthika G, Sharma N, Shahada C and Sudheer S (2023) Utilization of banana peel waste and its application: A review. *The Pharm InnoJ.*, 12(5): 2544 - 2548.
- Keeton JT (1983) Effect of fat and NaCl / phosphate levels on the chemical and sensory properties of pork patties. *J Food Sci.*, 48: 787- 885.
- Kim HY, Kim KJ, Lee JW, Kim GW and Kim CJ (2012) Effects of chicken feet gelatin and wheat fiber levels on quality properties of semi-dried chicken jerky. *Korean J. Food Sci Ani Reso.*, 32: 732–739.
- Kookal S. and Thimmaiah A (2018) Nutritional composition of staple food bananas of three cultivars in India. *American J Plant Sci.*, 9, 2480 - 2493.
- Lopez – Vargas JH, Fernandez – Lopez J, Perze – Alvaez JA and Viuda – Martos M (2014) Quality characteristics of pork burger added with albedo-fiber powder obtained from yellow passion fruit (*Passiflora edulis* var. *flavicarpa*) co-products. *Meat Sci.*, 97(2): 270 – 276.
- Ming-Chang Li , Chin-Fu Chou, Shu-Chen Hsu and Jen-Shinn Lin(2020)Physicochemical characteristics and resistant starch of different varieties of banana from Taiwan .*Int J Food Propert.*, 23 (1): 1168 – 1175.
- Murphy EW, Criner PE and Grey BC (1975) Comparison of methods for calculating retentions of nutrients in cooked foods. *J Agric Food Chem.*, 23:1153 – 1157.
- Petracci M, Bianchi M,Mudalal S and Cavani C (2013) Functional ingredients for poultry meat products, *Trends Food Sci. Technol.*, 33 (1): 27 - 39.
- Pereira J, Brohi SA, Malairaj S, Zhang W and Zhou G (2020) Quality of fat-reduced frankfurter formulated with unripe banana by products and pre-emulsified sunflower oil. *Int J Food Prop.*, 23(1): 420 - 433.
- Petracci M, Bianchi M, Mudalal S and Cavani C (2013) Functional ingredients for poultry meat products. *Trends Food Sci. Technol.*, 33(1): 27-39.
- Santhi DandKalaikannan A(2014) The effect of the addition of oat flour in low-fat chicken nuggets. *J. Nutr. Food Sci.* 4(1): 1-4.
- Salazar DM, Arancibia S Casado A, Viteri ME, Lopez-Caballero and Montero MP (2021). Green Banana (*Musa acuminata* AAA) wastes to develop an edible film for food applications. *Polymers* 13(18): 3183.
- Snedecor GW and Cochran WG 1968. *Statistical Methods*, 6thedn. Oxford and IBH Publishing Co. New Delhi. 299-338.
- Sultan Shaikh FR, Devangare AA, Raziuddin M, Gaikwad NZ, AwandkarSP, Suryawanshi RD and Kulkarni RC (2020) Effect of green banana pulp powder on microbiological quality of chevon nuggets during refrigerated storage. *J Meat Sci.*, 17 (2): 22 – 29.
- Taludkar S. and Sharma DP. (2010) Development of dietary fiber rich chicken meat patties using wheat and oat bran. *J. Food Sci. Technol.*, 47(2): 224–229.

- Trout ES, Hunt MC, Johnson DE, Claus JR, Kastner CL, Kropf DH and Stroda S (1992) Chemical, physical and sensory characterization of ground beef containing 5 to 30 percent fat. *J. Food Sci.*, 57: 25-29.
- Vu, HT, Scarlett CJ and Vuong QV (2018) Phenolic compounds within banana peel and their potential uses: A review. *J Funct Foods.*, 40: 238–248.
- Yadav S, Pathera AK, Malik A and Sharma DP (2020) Development of fibre enriched chicken nuggets using combination of wheat bran with dried apple pomace or dried carrot pomace. *J Ani Res.*, 10(!): 95-103.
- Zaini HBM, Mantihal S, Ng SWY and Pindi W (2021) The incorporation of green peas as the source of dietary fiber in developing functional chicken nuggets. *J Food Process Preserv.*, 00: e15412.